

Higgs Searches at Future Colliders: Potentials for Discovery and Diagnosis*

V. Barger^a

^aPhysics Department, University of Wisconsin, Madison, WI 53706

The prospects for Higgs boson discovery and study at present and future colliders are reviewed, with emphasis on the Minimal Supersymmetric Standard Model (MSSM) states. The expected experimental coverages with different production and decay modes are outlined. Particular attention is given to the Higgs couplings predicted when the top quark Yukawa coupling has an infrared fixed point. Characteristic Higgs boson mass spectra from the minimal supergravity (mSUGRA) model are considered and the decoupling/non-decoupling regimes of the lightest Higgs boson are discussed. General tests are described to ascertain whether a discovered light Higgs boson is a SM or MSSM state and to diagnose its fundamental properties. The constraints on the Higgs mass spectra implied by an interesting cosmological relic density of the lightest supersymmetric particle are described.

1. Introduction

“What breaks the electroweak symmetry?” is the central physics issue of our time. Higgs bosons are the cornerstones of our theoretical models. The crucial tests of the spontaneous symmetry breaking (EWSB) mechanism are the Higgs boson interactions with the W and Z gauge fields.

There are numerous options for Higgs multiplets. The Standard Model (SM) assumes one $SU(2)$ doublet of Higgs fields and has one physical Higgs boson (h_{SM}^0). The Minimal Supersymmetric Standard Model (MSSM) contains two doublets of Higgs fields and five physical Higgs bosons ($h^0, H^0, A^0, H^\pm$). Non-supersymmetric two-Higgs doublet models have a phase in the vacuum expectation values which is interesting for CP violation (e.g., a muon electric dipole moment). Higher dimensional Higgs multiplets such as $I = 3/2$ contain doubly charged Higgs fields. If no neutral Higgs bosons are found to exist with mass $m_h \lesssim 0.8$ TeV, then strong scattering must occur in the WW sector. The thrust of this review will be the SM and MSSM Higgs scenarios. The reader is referred to other comprehensive recent reviews ([1–4] and references therein) for more thorough discussions.

2. Indirect $m_{h_{\text{SM}}^0}$ determinations

Precision M_W and m_t measurements test radiative corrections through the relation

$$M_W = M_Z \left[1 - \frac{\pi\alpha}{\sqrt{2}G_\mu M_W^2 (1 - \delta r)} \right]^{1/2}. \quad (1)$$

In the SM the loop corrections δr depend on m_t^2 and $\log m_h$ (see e.g. Ref. [5]). The optimal relative precision for tests of this relation are

$$\Delta M_W \approx \frac{1}{140} \Delta m_t, \quad (2)$$

and hence the ΔM_W accuracy is most critical. The estimated achievable accuracies ΔM_W at future machines with integrated luminosities L are listed in Table 1. Note that a luminosity at the LHC higher than 10 fb^{-1} may not further improve ΔM_W . As Fig. 1 shows, precise ΔM_W measurement will give pinpoint accuracy on $m_{h_{\text{SM}}^0}$. A value $\Delta M_W = 6$ MeV (achievable at a muon collider with 100 fb^{-1} luminosity[6]) would determine $m_{h_{\text{SM}}^0}$ to

$$\Delta m_{h_{\text{SM}}^0} = \pm 10 \left(\frac{m_h}{100 \text{ GeV}} \right) \text{ GeV}. \quad (3)$$

New physics contributions to δr would likewise be stringently tested. An improved value of $\alpha(M_Z)$ from new measurements of $\sigma(e^+e^- \rightarrow \text{hadrons})$ at low energy is necessary to fully utilize a very accurate M_W measurement.

*Invited review talk at *The 5th International Conference on Supersymmetries in Physics (SUSY97)*, Philadelphia, Pennsylvania, May 27–31, 1997.

Table 1

Estimated accuracy of W -mass measurements at LEP-2, Tevatron, Large Hadron Collider, Next Linear Collider ($\sqrt{s} = 0.5$ TeV), and First Muon Collider ($\sqrt{s} = 0.5$ TeV). See Ref. [1].

	LEP-2		Tevatron		LHC	NLC (e^+e^-)	FMC ($\mu^+\mu^-$)
L (fb $^{-1}$)	0.1	2	2	10	10	50	50
ΔM_W (MeV)	144	34	35	20	15	15	9




Figure 1. Correlation between M_W and m_t in the SM with QCD and electroweak corrections for $m_h = 100, 300$ and 1000 GeV. The data point and error bars illustrate the possible accuracy for the indirect m_h determination assuming $M_W = 80.356 \pm 0.006$ GeV and $m_t = 175 \pm 0.2$ GeV. The widths of the bands indicate the uncertainty in $\alpha(M_Z)$. From Ref. [6].

3. SM Higgs – the benchmark

The Higgs mass $m_{h_{\text{SM}}^0}$ is the only parameter of the SM Higgs sector. SM Higgs production occurs through the processes of Fig. 2. QCD radiative corrections have been evaluated for all hadronic production processes but $t\bar{t}h$ and $b\bar{b}h$. The SM backgrounds to the Higgs signals are well studied.

The diagrams for the SM Higgs decays of primary interest are shown in Fig. 3. The partial widths for Higgs boson decays to fermions are proportional to m_f^2 , so a light SM Higgs decays dominantly to $b\bar{b}$, with $b\bar{b}$ branching fraction $\sim 90\%$. The rare $\gamma\gamma$ mode with branching frac-




Figure 2. Higgs boson production diagrams for (a) $\ell^+\ell^-$ and (b) $p\bar{p}, pp$ colliders.




Figure 3. Higgs boson decay diagrams

tion $\sim 10^{-3}$ for $m_{h_{\text{SM}}^0} \sim 80\text{--}140$ GeV is important for LHC searches. Sophisticated detectors are required with b -flavor tagging capabilities for $b\bar{b}$ decays and high EM calorimeter resolution ($\sim 1\%$) for $\gamma\gamma$ detection. For heavy SM Higgs searches via the mode $h \rightarrow ZZ \rightarrow 4\ell$ (with $\ell = e$ or μ), large angular coverage and muon detection are needed.

In the ongoing search at LEP-2 for the SM

Higgs, the collider energy is the major limiting factor. Figure 4 shows the potential 5σ discovery limits for $m_{h_{\text{SM}}^0}$ at $\sqrt{s} = 175, 192$ and 205 GeV versus the minimum luminosity needed per experiment (with results from 4 experiments combined). The ALEPH collaboration has placed the bound[7]

$$m_{h_{\text{SM}}} > 70.7 \text{ GeV} \quad (4)$$

from 21.3 pb^{-1} of luminosity in LEP-2 running up to the current energy $\sqrt{s} = 172$ GeV.




Figure 4. Minimum luminosity needed per experiment, in pb^{-1} , for a combined 5σ discovery (full line) or 95% exclusion (dashed line) as a function of Higgs boson mass, at three center-of-mass energies. From Ref. [3].

Upgrades of the Tevatron (Main Injector, TeV-33) will allow for a SM Higgs search up to $m_{h_{\text{SM}}} \sim 120 \text{ GeV}$ [8] and possibly higher[9], through the

Wh_{SM} production mode with $W \rightarrow \ell\nu$ and $h_{\text{SM}} \rightarrow b\bar{b}$ decays. The 5σ discovery requirements on luminosity for the detection are shown in Fig. 5.




Figure 5. Luminosity requirements for 5σ discovery of a SM Higgs boson at the Tevatron in Wh_{SM} production. From Ref. [8].

Table 2 summarizes the discovery reach for SM Higgs searches at present and future colliders. The coverage by various modes at the different colliders is illustrated graphically in Fig. 6. Note that the combined LEP-2 + Tevatron Main Injector + LHC searches will cover the entire interesting $m_{h_{\text{SM}}}$ range up to $\mathcal{O}(\text{TeV})$.

4. MSSM Higgs ($h^0, H^0, A^0, H^\pm$) – the target

In the supersymmetric scenario there is an upper bound on the mass of the lightest Higgs boson. At tree level this bound is (see e.g. Ref. [10])

$$m_{h^0} \leq M_Z |\cos 2\beta|, \quad (5)$$

Table 2

Discovery reach for SM Higgs bosons at present and future colliders.

Collider	Integrated Luminosity	$m_{h_{SM}}$ Discovery Reach (GeV)
LEP-2 $\sqrt{s} = 192$ GeV	150 pb $^{-1}$	95
LEP-2 $\sqrt{s} = 205$ GeV	150 pb $^{-1}$	102
Tevatron (M.I.)	10 fb $^{-1}$	80–100
TeV-33	30 fb $^{-1}$	120
LHC	100 fb $^{-1}$	800
NLC (e^+e^-), FMC ($\mu^+\mu^-$) $\sqrt{s} = 500$	50 fb $^{-1}$	350
NLC $\sqrt{s} = 1000$	200 fb $^{-1}$	800
NMC ($\mu^+\mu^-$) $\sqrt{s} = 4000$	200 fb $^{-1}$	TeV range




Figure 6. SM Higgs mass ranges over which statistically significant signals are expected in leading decay modes at future colliders (see Ref. [1]).

where $\tan \beta = v_u/v_d$ is the ratio of vacuum expectation values. Radiative corrections can substantially raise the tree level bound to [1–3]

$$m_{h^0} \lesssim 130 \text{ GeV}. \quad (6)$$

The value of m_{h^0} depends on $\tan \beta$, the stop masses and the stop mixing. In the large m_A limit the radiatively corrected m_h is given by [11]

$$m_h^2 = M_Z^2 \cos^2 2\beta \left(1 - \frac{3m_t^2 t}{8\pi^2 v^2} \right) + \frac{3m_t^4}{4\pi^2 v^2} \left[t + \frac{\tilde{\kappa}}{2} + \left(\frac{3}{32\pi^2} \frac{m_t^2}{v^2} - \frac{2\alpha_3}{\pi} \right) (\tilde{\kappa}t + t^2) \right] \quad (7)$$

where

$$t = \ln \left(\frac{M_s^2}{m_t^2} \right),$$

$$\kappa = \frac{2\tilde{A}_t^2}{M_S^2} \left(1 - \frac{\tilde{A}_t^2}{12M_S^2} \right),$$

$$\tilde{A}_t = A_t + \mu / \tan \beta,$$

$$M_S^2 \simeq m_{t_L}^2 + m_{t_R}^2.$$

In a non-minimal supersymmetric model with extra Higgs doublets and singlets, the mass of at least one Higgs boson satisfies [12]

$$m_{h^0} \lesssim 150 \text{ GeV} \quad (8)$$

so long as the couplings remain perturbative to the Planck scale.

The current LEP-2 lower bounds on the Higgs masses in the MSSM are [7]

$$m_{h^0} > 62.5 \text{ GeV}, \quad (9)$$

$$m_{A^0} > 62.5 \text{ GeV}, \quad (10)$$

$$m_{H^\pm} > 44 \text{ GeV}. \quad (11)$$

These limits are based on searches for the final states

$$H^0 Z(hZ) \rightarrow b\bar{b}q\bar{q}, b\bar{b}\ell\bar{\ell}, b\bar{b}\nu\bar{\nu}, \tau\bar{\tau}q\bar{q}, \quad (12)$$

$$hA \rightarrow b\bar{b}b\bar{b}, \tau\bar{\tau}b\bar{b}. \quad (13)$$

The production cross sections have the β -dependences

$$\sigma_{hZ}, \sigma_{HA} \propto \sin^2(\beta - \alpha), \quad (14)$$

$$\sigma_{HZ}, \sigma_{hA} \propto \cos^2(\beta - \alpha), \quad (15)$$

where α is the Higgs mixing angle. Figure 7 shows the regions of the $(\tan \beta, m_h)$ space presently excluded by ALEPH searches and theoretical constraints, with $M_S = 1$ TeV assumed.




Figure 7. The $(m_h, \tan\beta)$ plane in the maximal stop mixing configuration. The dark areas are theoretically disallowed. The hatched area is excluded at 95% confidence level by the combined search for $e^+e^- \rightarrow hZ$ and $e^+e^- \rightarrow hA$. The dot-dashed lines show the change in the theoretically excluded region in the no mixing configuration. From Ref. [7]

5. $\tan\beta$ from top Yukawa fixed point

In a unified MSSM a large top quark Yukawa coupling (λ_t) at the GUT scale is needed for radiative electroweak symmetry breaking and for b - τ Yukawa unification[13–15]. Then λ_t approaches an infrared point, giving a relation between m_t and $\tan\beta$ that is independent of the λ_t value at M_{GUT} . Figure 8 shows the λ_t fixed point regions in the $\tan\beta$ vs. m_t^{pole} plane. For $m_t = 175$ GeV there are two solutions

$$\tan\beta \approx 1.8 \quad (\lambda_b^{\text{GUT}} \simeq \lambda_\tau^{\text{GUT}}), \quad (16)$$

$$\tan\beta \approx 56 \quad (\lambda_t^{\text{GUT}} \simeq \lambda_b^{\text{GUT}} \simeq \lambda_\tau^{\text{GUT}}). \quad (17)$$

These fixed point solutions are very attractive theoretically, but it is premature to rule out $\tan\beta$ values between the fixed-point values. The Higgs couplings to fermions and to weak bosons are

strongly dependent on $\tan\beta$, as illustrated in Fig. 9. Search strategies must allow for large variations in the possible coupling strengths.




Figure 8. Contours of constant $m_b(m_b)$ in the $m_t(m_t), \tan\beta$ plane with contours of constant GUT scale Yukawa couplings. From Ref. [13].

6. mSUGRA model

The existence of a low-energy supersymmetry may be indicated by the successful unification of the gauge couplings. In minimal supergravity (mSUGRA) models, the Higgs boson mass spectrum depends mainly on the universal scalar (m_0) and gaugino ($m_{1/2}$) masses at the GUT scale and on $\tan\beta$ [16]. Figure 10 illustrates representative Higgs masses for the low and high $\tan\beta$ fixed point solutions[17]. For $\tan\beta \simeq 1.8$ the A^0 , H^0 , and $H^\pm$ are considerably more massive than h^0 , whereas for $\tan\beta \simeq 50$, the A^0 , H^0 , and $H^\pm$ can be comparable in mass to h^0 . In the low $\tan\beta$ fixed point scenario the h^0 has a mass range that is accessible to LEP-2.

There are two interesting h^0 coupling regimes, according to whether A is much larger or comparable in mass to the Z [18]; see Table 3.

At the LHC at least one of the MSSM Higgs bosons should be discovered. The coverage of the




Figure 9. MSSM couplings normalized to the SM couplings. From Ref. [3].

$(m_A, \tan \beta)$ plane from various channels is shown in Fig. 11, assuming $M_S = 1$ TeV and no squark mixing. However, for some regions of the parameter space only h could be found.

The discovery potential for MSSM Higgs boson at various colliders is summarized in Table 4. For complete coverage of the Higgs states, high energy e^+e^- and $\mu^+\mu^-$ colliders may be necessary if the masses of the heavier states are $\mathcal{O}(\text{TeV})$.

7. What we will learn from h searches

If m_h exceeds 130 GeV, the MSSM is inconsistent. If m_h exceeds 150 GeV, weakly coupled supersymmetry is inconsistent. If m_h is below 130 GeV, the MSSM is viable, but is the Higgs boson the MSSM Higgs? Discovery of only one Higgs boson is insufficient to establish the MSSM.

Table 3
Two h^0 coupling regimes.

decoupling	non-decoupling
$m_A \gg M_Z$	$m_A \approx \mathcal{O}(M_Z)$
$h^0 \sim h_{\text{SM}}^0$	$h^0 \neq h_{\text{SM}}^0$
h^0 searches same as h_{SM}^0	some h_{SM}^0 search strategies change
use precision measurements to show $h^0 \neq h_{\text{SM}}^0$ or discover $H^0, A^0, H^{\tilde{N}}$	$gg \rightarrow h$ dominates at LHC in SM and MSSM for $\tan \beta \lesssim 4$ but $gg \rightarrow b\bar{b}h^0$ dominates for $\tan \beta \gtrsim 10$

Table 4
Discovery potential for MSSM Higgs bosons

	complete coverage	partial coverage
LEP-2		h^0 $m_{h^0} \lesssim 95$ GeV $h^0, A^0, H^0, H^{\tilde{N}}$ possible
TeV-33		h^0 $m_{h^0} \lesssim 120$ GeV
LHC		at least one MSSM Higgs boson
NLC-500	h^0	$A^0, H^0, H^{\tilde{N}}$ $m_A < \frac{1}{2}\sqrt{s}$
NLC-500 ($\gamma\gamma$)	h^0	A^0, H^0 $m_A < 0.8\sqrt{s}$
FMC-500	h^0	A^0, H^0 $m_A < \sqrt{s}$




Figure 10. The neutralino relic density and SUSY Higgs mass spectrum versus $m_{1/2}$ for $\mu > 0$ with (a) $\tan\beta = 1.8$, $m_0 = 150$ GeV and (b) $\tan\beta = 50$, $m_0 = 600$ GeV. $\tilde{\nu}$ is the lightest scalar neutrino. The shaded regions denote the parts of the parameter space (i) producing $\Omega_{\chi_1} h^2 < 0.1$ or $\Omega_{\chi_1^0} h^2 > 0.5$, (ii) excluded by theoretical requirements, or (iii) excluded by the chargino search at LEP-2. From Ref. [17].

High energy lepton colliders would insure discovery of heavy Higgs bosons. At a 1.5 TeV e^+e^- collider, the $A, H^0, H^\pm$ states could be discovered for $m_A < 0.75$ TeV. At a 4 TeV $\mu^+\mu^-$ collider, these Higgs bosons can be found for $m_A < 4$ TeV.

8. h_{SM}^0 or h_{MSSM}^0 ?

There are two ways to know whether a Higgs boson is a SM or MSSM member. The first way is to discover the other MSSM Higgs bosons. The second way is to measure the h^0 branching fractions and total width. The enhanced couplings to $b\bar{b}$ and $\mu^+\mu^-$ of the h_{MSSM}^0 compared to h_{SM}^0 give MSSM/SM coupling ratios exceeding 1 by $\sim 10\%$ or more if $m_A \lesssim 400$ GeV. The h^0 branching fractions for the decays $h^0 \rightarrow b\bar{b}, c\bar{c}, ZZ^*, WW^*, \gamma\gamma$ can be measured to $\sim 10\%$ accuracy at e^+e^- , $\gamma\gamma$ and $\mu^+\mu^-$ colliders[1,2].

The total widths of the Higgs boson can be another sensitive gauge of the Higgs nature. For example, if $m_A \lesssim 150$ GeV, then $\Gamma_{\text{total}}(h_{\text{MSSM}}^0)$

grows proportional to $\tan^2\beta$; for $\tan\beta \sim 1.8$, $\Gamma_{\text{total}} \sim 5$ MeV but for $\tan\beta \sim 50$, $\Gamma_{\text{total}} \sim 0.5$ GeV. The s -channel production of Higgs bosons at a muon collider can be used to scan over the Breit-Wigner resonance profile and directly measure Γ_h [19].

9. Post-discovery: establish pedigree of Higgs boson

Once a Higgs boson is discovered the next step is to measure its properties. Is it a spin zero particle? If $h \rightarrow \gamma\gamma$ decay is observed then the spin $\neq 1$ by Yang's theorem. Its spin can be established by measuring angular distributions in $h \rightarrow b\bar{b}$, $h \rightarrow e^+e^-e^+e^-$, etc. Is the Higgs boson CP even, odd, or an admixture? The general forms of the couplings to fermions $\bar{f}(a + \kappa\gamma_5)f$ and Z bosons $g_{\mu\nu} + i\kappa\epsilon_{\mu\nu\alpha\beta}P_1^\alpha P_2^\beta/\Lambda^2$ can be measured from angular correlations[1]. Are the tree level coupling strengths proportional to m_f and M_Z^2 as predicted? Are the loop couplings $gg \rightarrow h$ and




Figure 11. MSSM Higgs discovery regions (5σ) in $(m_A, \tan\beta)$ parameter space for the ATLAS+CMS experiments at the LHC with 300 fb^{-1} per detector. From Ref. [20].

$h \rightarrow \gamma\gamma$ of the strengths predicted from known particles in the loops? The detailed studies required to establish the Higgs boson properties should be easier at e^+e^- and $\mu^+\mu^-$ colliders than at hadron colliders.

10. The dark matter card

With R -parity conservation, the lightest supersymmetric particle (LSP) is a natural candidate for the dark matter in the universe. In mSUGRA models the lightest neutralino χ_1^0 is the preferred LSP[16]. The LSP relic density $\Omega = \rho/\rho_{\text{critical}}$ is governed by the annihilation cross section of LSPs in the early universe[21] and the $\chi_1^0\chi_1^0 \rightarrow \text{Higgs and } Z \text{ resonance poles}$ are very influential[17,22]. Hence the relic density depends on the masses and widths of the neutral Higgs bosons. A cosmologically interesting relic density

$$0.1 \lesssim \Omega h^2 \lesssim 0.5 \quad (18)$$

implies significant constraints on the mSUGRA parameters and the Higgs mass spectrum, as shown in Fig. 10. As the cold dark matter component of Ωh^2 becomes more precisely known over the next few years, the corresponding restrictions on the Higgs mass spectrum will be increasingly interesting.

11. Conclusions

Electroweak symmetry breaking and mass generation are mysteries that will be solved by forthcoming collider experiments. Precision electroweak tests already point the smoking gun towards Higgs bosons. It is still a horse race to the discovery line for h^0 . At LEP-2 the energy is critical. At the Tevatron the luminosity is critical. At the LHC at least one Higgs boson should be found. The genetic traits (couplings, branching fractions, total widths) of the lightest Higgs are critical clues to decide if it is the SM, MSSM, a generic 2-doublet member, or otherwise. The top Yukawa infrared fixed points at $\tan\beta \approx 1.8$ and $\tan\beta \approx 56$ in the unified MSSM are particularly attractive scenarios; the Higgs boson couplings are very different at low and high $\tan\beta$. Decoupling of the light Higgs boson ($h^0 \approx h_{\text{SM}}^0$) occurs at large m_A ; for non-decoupling all MSSM Higgs bosons are relatively light. The Next Linear Collider (e^+e^-) and First Muon Collider ($\mu^+\mu^-$) are essential for definitive studies of Higgs boson properties (spin, CP, decay widths). The high energy lepton colliders NLC(e^+e^-)-1.5 TeV and NMC($\mu^+\mu^-$)-4 TeV would provide the energy reach for a high mass Higgs spectrum. Finally, the existence of an interesting cosmological relic density imposes strong constraints on the mSUGRA parameter space and the Higgs boson masses.

Acknowledgments: I thank Chung Kao and Tao Han for valuable advice in the preparation of this talk. This work was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-95ER40896 and in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation.

REFERENCES

1. H. Haber, T. Han, F.S. Merritt, J. Womersley et al., hep-ph/9703391, in Proceedings of the 1996 DPF/DPB Summer Study on New Directions for High Energy Physics, Snowmass, CO; H. Haber, Ringberg Workshop on the Higgs Puzzle, Ringberg Castle, Germany (1996), hep-ph/9703381.
2. J.F. Gunion, L. Poggioli, R. Van Kooten, C. Kao, P. Rowson et al., hep-ph/9703330, in proceedings of Ref. 1.
3. M. Carena et al., hep-ph/9602250, in Vol. 1, Report of the Workshop on Physics at LEP2, G. Altarelli, T. Sjostrand and F. Zwirner (eds), CERN 96-01.
4. S. Dawson, hep-ph/9703387, to be published in *Perspectives in Higgs Physics*, ed. by G. Kane.
5. P. Langacker and J. Erler, in Review of Particle Physics, Phys. Rev. **D54**, 85 (1996); K. Hagiwara, D. Haidt, and S. Matsumoto, hep-ph/9706331 (1997).
6. V. Barger, M.S. Berger, J.F. Gunion, and T. Han, Univ. of Wisconsin-Madison preprint MADPH-96-963 [hep-ph/9702334], to appear in Phys. Rev. **D56** (1997).
7. The ALEPH Collaboration, CERN reports CERN-PPE/97-070 and CERN-PPE-97-071 (1997); S.L. Wu (private communication).
8. S. Kim, S. Kuhlmann, and W.M. Yao, CDF/ANAL/EXOTIC/PUBLIC/3904, in Proceedings of the 1996 DPF/DPB Summer Study on New Directions for High-Energy Physics, Snowmass, CO.
9. A. Stange, W. Marciano, and S. Willenbrock, Phys. Rev. **D49**, 1354 (1994); **D50**, 4491 (1992); S. Mrenna and G.L. Kane, hep-ph/9406337 (1994).
10. J.F. Gunion, H.E. Haber, G.L. Kane, and S. Dawson, *The Higgs Hunters Guide*, (Addison-Wesley, 1990).
11. M. Carena, J.R. Espinosa, M. Quiros, and C.E.M. Wagner, Phys. Lett. **355B**, 209 (1995).
12. C. Kolda, G. Kane, and J. Wells, Phys. Rev. Lett. **70**, 2686 (1993).
13. V. Barger, M.S. Berger, and P. Ohmann, Phys. Rev. **D47**, 1093 (1993); V. Barger, M.S. Berger, and P. Ohmann, in *Proceedings of the XXVII Int. Conf. on High Energy Physics*, Glasgow, July 1994, (IOP Publishing Ltd., UK, 1995), p. 563.
14. V. Barger, M.S. Berger, P. Ohmann, and R.J.N. Phillips, Phys. Lett. **B314**, 3511 (1993).
15. J. Bagger, S. Dimopoulos, and E. Masso, Phys. Rev. Lett. **55**, 920 (1985); H. Arason et al., Phys. Rev. Lett. **67**, 2933 (1991); Phys. Rev. **D46**, 3945 (1992); C.D. Froggatt, R.G. Moorhouse, and I.G. Knowles, Phys. Lett. **B298**, 356 (1993); P. Langacker and N. Polonsky, Phys. Rev. **D50**, 2199 (1994); W.A. Bardeen, M. Carena, S. Pokorski, and C.E.M. Wagner, Phys. Lett. **B320**, 110 (1994); M. Carena, M. Olechowski, S. Pokorski, and C.E.M. Wagner, Nucl. Phys. **B452**, 45 (1995); B. Schrempp, Phys. Lett. **B344**, 193 (1995); B. Schrempp and M. Wimmer, DESY-96-109 (1996), hep-ph/9606386, and references therein.
16. J. Ellis and F. Zwirner, Nucl. Phys. **B338**, 317 (1990); G. Ross and R.G. Roberts, Nucl. Phys. **B377**, 571 (1992); R. Arnowitt and P. Nath, Phys. Rev. Lett. **69**, 725 (1992); M. Drees and M. Nojiri, Nucl. Phys. **B369**, 54 (1993); S. Kelley et al., Nucl. Phys. **B398**, 3 (1993); M. Olechowski and S. Pokorski, Nucl. Phys. **B404**, 590 (1993); V. Barger, M.S. Berger, and P. Ohmann, Phys. Rev. **D49**, 4908 (1994); G. Kane, C. Kolda, L. Roszkowski, and J. Wells, Phys. Rev. **D49**, 6173 (1994); D.J. Castaño, E. Piard and P. Ramond, Phys. Rev. **D49**, 4882 (1994); W. de Boer, R. Ehret and D. Kazakov, Z. Phys. **67**, 647 (1995); H. Baer, M. Drees, C. Kao, M. Nojiri, and X. Tata, Phys. Rev. **D50**, 2148 (1994); H. Baer, C.-H. Chen, R. Munroe, F. Paige, and X. Tata, Phys. Rev. **D51**, 1046 (1995).
17. V. Barger and C. Kao, UW-Madison report no. MADPH-97-992, hep-ph/9704403 (1997).
18. H.E. Haber and Y. Nir, Nucl. Phys. **B335**, 363 (1990).
19. V. Barger, M.S. Berger, J. Gunion, and T. Han, Phys. Rep. **286**, no. 1, p. 1 (1997).

20. E. Richter-Was, D. Froidevaux, F. Gianotti, L. Poggioli, D. Cavalli, S. Resconi, CERN-TH-96-111 (1996).
21. B. Lee and S. Weinberg, Phys. Rev. Lett. **39**, 165 (1977); R. Barbieri, M. Frigeni, and G.F. Guidice, Nucl. Phys. **B313**, 725 (1989); J.L. Lopez, D.V. Nanopoulos, and K.-J. Yuan, Phys. Lett. **B267**, 219 (1991); Nucl. Phys. **B370**, 445 (1992); J. Ellis and L. Roszkowski, Phys. Lett. **B283**, 252 (1992); L. Roszkowski and R. Roberts, Phys. Lett. **B309**, 329 (1993); M. Kawasaki and S. Mizuta, Phys. Rev. **D46**, 1634 (1992); G.L. Kane, C. Kolda, L. Roszkowski, and J.D. Wells, Phys. Rev. **D49**, 6173 (1994); E. Diehl, G.L. Kane, C. Kolda, and J.D. Wells, Phys. Rev. **D52**, 4223 (1995); M. Drees and M. Nojiri, Phys. Rev. **DD47**, 376 (1993); R. Arnowitt and P. Nath, Phys. Lett. **B299**, 58 (1993); **B307**, 403(E) (1993); Phys. Rev. Lett. **70**, 3696 (1993); M. Drees and A. Yamada, Phys. Rev. **D53**, 1586 (1996); G. Jungman, M. Kamionkowski, and K. Griest, Phys. Rep. **267** (1996).
22. K. Griest and D. Seckel, Phys. Rev. **D43**, 3191 (1991); P. Goldono and G. Gelmini, Nucl. Phys. **B360**, 145 (1991); H. Baer and M. Brhlik, Phys. Rev. **D53**, 597 (1996); J. Ellis, T. Falk, K.A. Olive, and M. Schmitt, Phys. Lett. **B388**, 97 (1996); CERN Report no. CERN-TH-97-105, hep-ph/9705444 (1997).